

Stability of Refrigerated and Frozen Drugs

—Chart modified November 2008—

(Based on U.S. product labeling and relevant studies)

Failure to follow storage recommendations of pharmaceutical products can result in subpotent products and potentially, therapeutic failure.¹⁻³ In addition, hundreds of thousand of medications are discarded each year because of improper storage conditions resulting in millions of dollars wasted.^{1,2} This can occur for a variety of reasons including improper shipping, failure to refrigerate or freeze a product upon receipt or after use, and in the event of a power failure. It is important to recognize when pharmaceutical products have not been stored according to the manufacturer's specifications and take appropriate action. Depending on the product, lot number and expiration date, time of exposure to temperatures outside of the recommended range and actual temperature of exposure, some medications may be deemed suitable for administration while others should be discarded. The following chart provides general guidelines for the room temperature stability of common medications and vaccines which are stored in the refrigerator or freezer. However, if a pharmaceutical product has not been stored according to the recommended conditions, it is important to contact the manufacturer and provide the specifics (lot number, expiration date, exposure temperature, length of time) to determine if the product is viable. Continue to the end of this chart for helpful tips on what to do during a power failure or when there is a potential for flooding.

• For a listing of Anti-infective Suspensions and their stability after reconstitution, see *Detail-Document #231107*

• For a listing of Insulin Products and their storage/stability information, see *Detail-Document #220910*

Generic name (<i>Brand Name</i>)*	Recommended Storage**	Stability at Room Temperature** FOR SPECIFIC INFORMATION, CONTACT MANUFACTURER	Manufacturer Contact Information (Quality Control)
Adalimumab (<i>Humira</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Abbott Laboratories 800-633-9110
Alprostadil injection (<i>Caverject solution</i>)	Refrigerate 40 mcg strength at 36° to 46° F (2° to 8°C) until dispensed. (Store the 5, 10, and 20 mcg strengths at or below 77°F [25°C]).	After dispensing 40 mcg strength: Three months at or below 25°C (77°F).	Pfizer 800-438-1935
Alprostadil urethral suppository (<i>Muse</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	14 days.	Vivus, Inc. 650-934-5200

More . . .

Generic name (<i>Brand Name</i>)*	Recommended Storage**	Stability at Room Temperature** FOR SPECIFIC INFORMATION, CONTACT MANUFACTURER	Manufacturer Contact Information (Quality Control)
Amphotericin B (<i>Amphocin</i> , others)	Refrigerate at 36° to 46°F (2° to 8°C) before reconstitution (lyophilized powder).. After reconstitution, refrigerated solution (5 mg/mL concentration, diluted with Sterile Water for Injection) stable for one week.	24 hours (reconstituted solution at concentration of 5 mg/mL, diluted with Sterile Water for Injection).	Pfizer 800-438-1935
Azithromycin ophthalmic (<i>Azasite</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	Once the bottle is opened, store at 2° to 25°C (36° to 77°F) for up to 14 days. Discard after 14 days.	Inspire Pharmaceuticals 888-881-4696
Becaplermin (<i>Regranex</i>)	Refrigerate at 36° to 46°F (2° to 8°C). Do not freeze.	Contact manufacturer.*** Off-label [§] information indicates that unopened bottles left out of refrigerator are stable (up to 86°F or 30°C) for up to six days or less, one time only. ¹⁴	Johnson and Johnson Wound Management 888-REGRANEX
Benzoyl peroxide/clindamycin (<i>Duac</i>)	Refrigerate at 36° to 46°F (2° to 8°C). Do not freeze.	60 days.	Stiefel Labs 888-500-3376
Calcitonin nasal (<i>Fortical</i>)	Refrigerate unopened bottle at 36° to 46° F (2° to 8°C).	After opening, store bottle in use in an upright position for up to 30 days at 68° to 77°F (20° to 25°C). Excursions permitted to 59°F to 86°F (15° to 30°C). Off-label [§] information indicates that unopened bottles left out of refrigerator (up to 77°F or 25°C) in the pharmacy for seven days or less can be safely dispensed to the patient. ⁴ Contact manufacturer.***	Upsher-Smith Laboratories 763-315-2000

Generic name (<i>Brand Name</i>)*	Recommended Storage**	Stability at Room Temperature** FOR SPECIFIC INFORMATION, CONTACT MANUFACTURER	Manufacturer Contact Information (Quality Control)
Calcitonin injectable (<i>Miacalcin</i>)	Refrigerate unopened bottle at 36° to 46° F (2° to 8°C).	Contact manufacturer.***	Novartis 888-669-6682
Calcitonin nasal (<i>Miacalcin</i>)	Refrigerate unopened bottle at 36° to 46° F (2° to 8°C).	Opened bottle may be stored at 59°F to 86°F (15° to 30°C) in upright position for 35 days.	Novartis 888-669-6682
Chlorambucil (<i>Leukeran</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	Contact manufacturer.*** Off-label [§] information indicates stable when maintained at up to 86°F (30°C) for up to one week. ⁵	GlaxoSmithKline 866-475-8222
Darbepoetin alfa (<i>Aranesp</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Amgen 800-772-6436
Desmopressin (<i>DDAVP Rhinal Tube</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	Up to three weeks.	Sanofi Aventis 908-243-6000
Dinoprostone cervical gel (<i>Prepidil</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	Contact manufacturer.*** Off-label [§] information indicates stable when maintained for 24 hours. ⁶	Pfizer 800-438-1935
Dornase alfa (<i>Pulmozyme</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	Up to 24 hours. Contact manufacturer.*** Off-label [§] information indicates stable when maintained for up to 48 cumulative hours at a temperature up to 86°F (30°C). ⁷	Genentech 650-225-1000

Generic name (<i>Brand Name</i>)*	Recommended Storage**	Stability at Room Temperature** FOR SPECIFIC INFORMATION, CONTACT MANUFACTURER	Manufacturer Contact Information (Quality Control)
Dronabinol (<i>Marinol</i>)	Refrigerate at 36° to 46° F (2° to 8°C) or store in a cool environment between 46°F to 59°F (8°C to 15°C).	Contact manufacturer.*** Off-label [§] information indicates stable when maintained at room temperature 77°F (25°C) for 24 months. [§]	Solvay Pharmaceuticals 770-578-9000
Epoetin alfa (<i>Epogen</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Amgen 800-772-6436
Epoetin alfa (<i>Procrit</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.*** Off-label [§] information indicates the following stability data when maintained at room temperature up to 86°F (30°C): Single-dose vial – 14 days. ¹⁵ Unopened multi-dose vial – seven days. ¹⁵	Ortho Biotech 888-227-5624
Estradiol/norethindrone acetate transdermal system (<i>CombiPatch</i>)	Prior to dispensing, refrigerate at 36° to 46°F (2° to 8°C).	After dispensing, store at room temperature below 77°F (25°C) for up to six months.	Novartis 888-669-6682
Etonogestrel/ethinyl estradiol vaginal ring (<i>NuvaRing</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	After dispensing, store at 77°F (25°C) for up to four months; excursions permitted to 59° to 86°F (15° to 30°C).	Organon 877-688-2746
Estramustine (<i>Emcyt</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Pfizer 800-438-1935
Etanercept injection (<i>Enbrel</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.*** However, one reference suggests: ¹ Vial – seven days. ¹ Prefilled syringe – four days. ^{1,16}	Immunex Corporation 888-436-2735
Etoposide capsules (<i>VePesid</i>)	Prior to dispensing, refrigerate at 36° to 46° F (2° to 8°C).	Contact manufacturer.***	Bristol-Myers Squibb 609-897-2000

Generic name (<i>Brand Name</i>)*	Recommended Storage**	Stability at Room Temperature** FOR SPECIFIC INFORMATION, CONTACT MANUFACTURER	Manufacturer Contact Information (Quality Control)
Exenatide (<i>Byetta</i>)	Prior to first use, refrigerate at 36° to 46° F (2° to 8°C). Do not freeze.	After first use, store at a room temperature not to exceed 77°F (25°C). Discard after 30 days.	Amylin Pharmaceuticals 800-349-8919
Filgrastim (<i>Neupogen</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	Stable at room temperature for up to 24 hours.	Amgen 800-772-6436
Formoterol (<i>Foradil</i>)	Prior to dispensing, refrigerate at 36° to 46° F (2° to 8°C).	Store at room temperature 68° to 77°F (20° to 25°C). Off-label [§] information indicates stable when maintained at room temperature for up to 6 months. ⁹	Schering Corporation 800-222-7579
Gabapentin oral suspension (<i>Neurontin</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	Contact manufacturer.*** Off-label [§] information indicates oral solution stable for seven days at temps up to 86°F (30°C). ⁶	Pfizer 800-438-1935
Glatiramer acetate injection (<i>Copaxone</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	One month at room temperature of 59° to 86°F (15° to 30°C).	Teva Neuroscience 888-838-2872
Interferon alfacon-1 (<i>Infergen</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Three Rivers Pharm. 724-778-6100
Interferon beta-1a (<i>Avonex</i>)	Refrigerate unconstituted vial of lyophilized powder at 36° to 46°F (2° to 8°C).	Lyophilized powder vial – 30 days. Prefilled syringes – seven days.	Biogen Idec 800-456-2255
Interferon beta-1a (<i>Rebif</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	30 days.	Pfizer 800-438-1935

Generic name (<i>Brand Name</i>)*	Recommended Storage**	Stability at Room Temperature** FOR SPECIFIC INFORMATION, CONTACT MANUFACTURER	Manufacturer Contact Information (Quality Control)
Interferon gamma-1b (<i>Actimmune</i>)	Refrigerate at 36° to 46°F (2° to 8°C) immediately upon receipt to ensure optimal retention of physical and biochemical integrity. Do not freeze.	Unopened vial should not be left at room temperature for a total time exceeding 12 hours prior to use. Vials stored at room temperature should not be returned to the refrigerator.	InterMune, Inc. 888-486-6411
Latanoprost (<i>Xalatan</i>)	Prior to dispensing and opening, store unopened bottles in refrigerator at 36°F to 46°F (2°C to 8°C).	During shipment, unopened bottle stable at up to 104°F (40°C) for no more than eight days. After first use, store at a room temperature not to exceed 77°F (25°C) for up to 6 weeks. Off-label [§] information indicates unopened bottle stable when maintained at continuous room temperature 77°F (25°C) for 12 months. ⁶ Contact manufacturer.***	Pfizer 800-438-1935
Liotrix (<i>Thyrolar</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	Contact manufacturer.***	Forest Pharmaceuticals 1-800-678-1605, ext. 66297
Lopinavir/ritonavir solution (<i>Kaletra</i>)	Refrigerate at 36° to 46°F (2° to 8°C) until dispensed. Stable until expiration date if stored in refrigerator.	Two months.	Abbott Laboratories 800-633-9110
Lorazepam intensol oral concentrate	Refrigerate at 36° to 46° F (2° to 8°C).	Contact manufacturer.*** Off-label [§] information indicates stable when maintained at continuous room temperature 77°F (25°C) for 30 days. ¹⁰	Roxane Laboratories 614-276-4000

Generic name (<i>Brand Name</i>)*	Recommended Storage**	Stability at Room Temperature** FOR SPECIFIC INFORMATION, CONTACT MANUFACTURER	Manufacturer Contact Information (Quality Control)
Melphalan tablets (<i>Alkeran</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	Contact manufacturer.***	GlaxoSmithKline 888-825-5249
Octreotide (<i>Sandostatin</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	At room temperature (70°F to 86°F, 20°C to 30°C), stable for 14 days, when protected from light.	Novartis 888-669-6682
Octreotide injectable suspension (<i>Sandostatin LAR</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	Drug kit can remain at room temperature for 30 to 60 minutes prior to preparation. However, must be used immediately after preparation.	Novartis 888-669-6682
Papain/urea topical (<i>Ethezyme</i>) PRODUCT DISCONTINUED BY MANUFACTURER	Refrigerate at 36° to 46°F (2° to 8°C) or in cool (46° to 59°F) place (8° to 15°C).	Contact manufacturer.*** Off-label [§] information indicates stable when maintained up to 77°F (25°C) for three months. ¹¹	Ethex 800-321-1705
Peginterferon alfa-2a (<i>Pegasys</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	24 hours.	Roche Laboratories 800-526-0625
Peginterferon alfa-2b (<i>PegIntron</i>)	<i>PegIntron</i> vials - Store at room temperature (77°F [25°C]; excursion permitted to 59°F to 86°F [15°C to 30°C]). After reconstitution, the solution should be used immediately, but may be stored up to 24 hours at 36° to 46°F (2° to 8°C). <i>PegIntron Redipen</i> - Refrigerate at 36° to 46° F (2° to 8°C). After reconstitution, the solution should be used immediately, but may be stored up to 24 hours at 36° to 46°F (2° to 8°C).	Contact manufacturer.*** Off-label [§] information indicates the <i>PegIntron Redipen</i> (unreconstituted lyophilized powder) is stable when maintained up to 86°F (30°C) for one day. However, there is no data to address the re-refrigeration of the product after maintained in an un-refrigerated state. ⁹	Schering Corporation 800-222-7579

Generic name (<i>Brand Name</i>)*	Recommended Storage**	Stability at Room Temperature** FOR SPECIFIC INFORMATION, CONTACT MANUFACTURER	Manufacturer Contact Information (Quality Control)
Pramlintide injection (<i>Symlin</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	30 days.	Amylin Pharmaceuticals 800-349-8919
Promethazine suppositories (<i>Promethegan</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	G & W Laboratories 908-753-2000
Proparacaine ophthalmic solution	Refrigerate at 36° to 46° F (2° to 8°C).	Contact manufacturer.***	Akorn 847-279-6100
Ritonavir capsules (<i>Norvir</i>)	Refrigerate at 36° to 46°F (2° to 8°C) until dispensed. Stable until expiration date if stored in refrigerator.	30 days.	Abbott Laboratories 800-633-9110
Sinecatechins (<i>Veregen</i>)	Prior to dispensing to the patient, refrigerate at 36° to 46°F (2° to 8°C).	After dispensing, can be stored at room temperature.	Doak Dermatologics 800-929-9300
Sirolimus solution (<i>Rapamune</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Up to 15 days.	Wyeth Pharmaceuticals 800-395-9938
Teriparatide (<i>Forteo</i>)	Refrigerate at 36° to 46° F (2° to 8°C). Minimize time out of refrigerator during use.	Contact manufacturer.***	Eli Lilly 800-545-5979
Tobramycin for oral inhalation (<i>Tobi</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	Opened or unopened pouches may be stored at room temperature (up to 77°F or 25°C) for up to 28 days.	Novartis 888-669-6682
Trifluridine (<i>Viroptic</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	Contact manufacturer.*** Off-label [§] information indicates stable when maintained up to 77°F (25°C) for two weeks. However, due to slight changes in pH (no change in potency), ocular stinging or discomfort may occur. ¹⁷	King Pharmaceuticals 800-776-3637

Generic name (<i>Brand Name</i>)*	Recommended Storage**	Stability at Room Temperature** FOR SPECIFIC INFORMATION, CONTACT MANUFACTURER	Manufacturer Contact Information (Quality Control)
Vaccines			
Diphtheria, tetanus toxoids (DT) (pediatric) Tetanus, diphtheria toxoids (Td) (adult) (<i>Decavac</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Sanofi Pasteur 800-822-2463
Diphtheria, tetanus toxoids, acellular pertussis vaccine (DTaP) (pediatric) (<i>Tripedia, Daptacel</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Sanofi Pasteur 800-822-2463
Diphtheria, tetanus toxoids, acellular pertussis vaccine (DTaP) (pediatric) (<i>Infanrix</i>) Diphtheria, tetanus toxoids, acellular pertussis vaccine, hepatitis B vaccine, inactivated polio vaccine (DTaP/HepB/IPV) (pediatric) (<i>Pediarix</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.*** Off-label [§] information indicates stable when maintained above 46°F to 77°F (above 8°C to 25°C) for 72 hours (<i>Infanrix</i>). ¹² Contact manufacturer.*** Off-label [§] information indicates stable when maintained above 46°F to 77°F (above 8°C to 25°C) for 72 hours (<i>Pediarix</i>). ¹²	GlaxoSmithKline 866-475-8222

Generic name (<i>Brand Name</i>)*	Recommended Storage**	Stability at Room Temperature** FOR SPECIFIC INFORMATION, CONTACT MANUFACTURER	Manufacturer Contact Information (Quality Control)
Tetanus toxoid, reduced, diphtheria toxoids, acellular pertussis vaccine (Tdap) (adult) (<i>Boostrix</i>)	Refrigerate at 36° to 46°F (2° to 8°C)	Contact manufacturer.*** Off-label [§] information indicates stable when maintained above 46°F to 77°F (above 8°C to 25°C) for 72 hours (<i>Boostrix</i>). ¹²	GlaxoSmithKline 866-475-8222
Diphtheria and Tetanus Toxoids and Acellular Pertussis Adsorbed, Inactivated Poliovirus and Haemophilus b Conjugate (Tetanus Toxoid Conjugate) Vaccine (pediatric) (<i>Pentacel</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Sanofi Pasteur 800-822-2463
<i>Haemophilus influenzae</i> type B conjugate vaccine (Hib) (<i>PedvaxHIB</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Merck 800-637-2590
<i>Haemophilus influenzae</i> type B conjugate vaccine (Hib) (<i>ActHIB</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Sanofi Pasteur 800-822-2463
<i>Haemophilus influenzae</i> type B conjugate (Hib) and hepatitis B vaccines (<i>Comvax</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Merck 800-637-2590
Hepatitis A vaccine (<i>Havrix</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.*** Off-label [§] information indicates stable when maintained above 46°F to 77°F (above 8°C to 25°C) for 72 hours. ¹²	GlaxoSmithKline 866-475-8222

Generic name (<i>Brand Name</i>)*	Recommended Storage**	Stability at Room Temperature** FOR SPECIFIC INFORMATION, CONTACT MANUFACTURER	Manufacturer Contact Information (Quality Control)
Hepatitis A vaccine (<i>Vaqta</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Merck 800-637-2590
Hepatitis A and Hepatitis B vaccine (<i>Twinrix</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	Contact manufacturer.*** Off-label information indicates stable when maintained above 46°F to 77°F (above 8°C to 25°C) for 72 hours (<i>Twinrix</i>). ¹²	GlaxoSmithKline 866-475-8222
Hepatitis B vaccine (<i>Engerix-B</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.*** Off-label [§] information indicates stable when maintained above 46°F to 77°F (above 8°C to 25°C) for 72 hours. ¹²	GlaxoSmithKline 866-475-8222
Hepatitis B vaccine (<i>Recombivax HB</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Merck 800-637-2590
Human papillomavirus (Types 6, 11, 16, 18) vaccine (<i>Gardasil</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Merck 800-637-2590
Inactivated polio vaccine (IPV) (<i>IPOL</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Sanofi Pasteur 800-822-2463
Influenza virus vaccine, types A and B (<i>Afluria</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	CSL Limited 888-4FLU-OFF

Generic name (<i>Brand Name</i>)*	Recommended Storage**	Stability at Room Temperature** FOR SPECIFIC INFORMATION, CONTACT MANUFACTURER	Manufacturer Contact Information (Quality Control)
Influenza virus vaccine, types A and B (<i>FluLaval</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	GlaxoSmithKline 866-475-8222
Influenza virus vaccine, types A and B (<i>Fluarix</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	GlaxoSmithKline 866-475-8222
Influenza virus vaccine, types A and B (<i>Fluvirin</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Novartis 800-244-7668
Influenza virus vaccine, types A and B (<i>Fluzone</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Sanofi Pasteur 800-822-2463
Influenza attenuated influenza vaccine, live (LAIV) (<i>FluMist</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.*** Off-label [§] information indicates stable when maintained above 46°F to 77°F (above 8°C to 25°C) for 8 hours. ¹³	MedImmune 800-949-3789
Measles, mumps, rubella vaccine (MMR) (<i>M-M-R II</i>)	Refrigerate at 36° to 46° F (2° to 8°C). Diluent may be stored in the refrigerator or at room temperature. Do not freeze diluent.	Contact manufacturer.***	Merck 800-637-2590

Generic name (<i>Brand Name</i>)*	Recommended Storage**	Stability at Room Temperature** FOR SPECIFIC INFORMATION, CONTACT MANUFACTURER	Manufacturer Contact Information (Quality Control)
Measles, mumps, rubella (MMR) and varicella vaccines (<i>ProQuad</i>)	Store in freezer at an average temperature of 5°F (-15°C) or colder for up to 18 months. Can be stored in a refrigerator at 36° to 46°F (2° to 8°C) for up to 72 hours before reconstitution. Diluent may be stored in the refrigerator or at room temperature. Do not freeze diluent.	Contact manufacturer.***	Merck 800-637-2590
Meningococcal conjugate, groups A, C, Y and W-135 vaccine (MCV) (<i>Menactra</i> [also contains diphtheria toxoid], <i>Menomune</i>)	Refrigerate at 35° to 46°F (2° to 8°C). (<i>Menomune</i> single-dose vial should be used within 30 minutes of reconstitution)	Contact manufacturer.***	Sanofi Pasteur 800-822-2463
Pneumococcal conjugate vaccine (<i>Prennar</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Wyeth 800-999-9384
Pneumococcal polysaccharide vaccine (<i>Pneumovax 23</i>)	Refrigerate at 36° to 46°F (2° to 8°C).	Contact manufacturer.***	Merck 800-637-2590
Rotavirus vaccine (<i>Rotarix</i>)	Refrigerate at 36° to 46°F (2° to 8°C). Diluent may be stored at room temperature.	After reconstitution, can be stored at room temperature up to 77°F (25°C) for 24 hours.	GlaxoSmithKline 866-475-8222
Rotavirus vaccine (<i>RotaTeq</i>)	Refrigerate at 36° to 46°F (2° to 8°C). Administer as soon as possible after removal from refrigerator.	Contact manufacturer.***	Merck 800-637-2590

Generic name (<i>Brand Name</i>)*	Recommended Storage**	Stability at Room Temperature** FOR SPECIFIC INFORMATION, CONTACT MANUFACTURER	Manufacturer Contact Information (Quality Control)
Varicella (chickenpox) vaccine (<i>Varivax</i>)	Store in freezer at 5° F (-15° C) or colder. Diluent may be stored in the refrigerator or at room temperature. Do not freeze diluent.	May be stored at refrigerator temperatures, 36° to 46° F (2° to 8°C) for up to 72 hours prior to reconstitution. Vaccine stored in refrigerator must be used within 72 hours of removal from freezer. Other storage conditions are determined on a case-by-case basis. Contact manufacturer.***	Merck 800-9-VARIVAX
Typhoid vaccine, live oral (<i>Vivotif</i>)	Refrigerate at 36° to 46° F (2° to 8°C).	Contact manufacturer.***	Berna Biotech, Ltd. 305-443-2900
Zoster (shingles) vaccine (<i>Zostavax</i>)	Store in freezer at 5° F (-15° C) or colder. Diluent may be stored in the refrigerator or at room temperature. Do not freeze diluent.	Contact manufacturer.***	Merck 800-9-VARIVAX

* This table includes commonly used medications and vaccines. For those products not listed, consult the product information or pharmaceutical manufacturer for stability information at room temperature.

** Data based on information provided by prescribing information, unless otherwise indicated. The following PIs were used for this chart: *Humira* (January 2003), *Caverject* (October 2003), *Muse* (August 2003), *Amphocin* (October 2006), *Azasite* (December 2007), *Regranex* (May 2008), *Duac* (August 2002), *Fortical* (February 2004), *Miacalcin* injection (November 2002), *Miacalcin* nasal spray (June 2006), *Leukeran* (November 2006), *Aranesp* (August 2008), *DDAVP Rhinal Tube* (October 2007), *Prepidil* (July 2008), *Pulmozyme* (2008), *Marinol* (March 2008), *Epogen* (August 2008), *Procrit* (August 2008), *CombiPatch* (January 2005), *NuvaRing* (December 2007), *Emcyt* (June 2007), *Enbrel* (June 2008), *VePesid* (July 2006), *Byetta* (June 2008), *Neupogen* (September 2007), *Foradil* (June 2006), *Neurontin* oral solution (January 2007), *Copaxone* (November 2007), *Infergen* (July 2006), *Avonex* (November 2006), *Rebif* (April 2008), *Actimmune* (January 2007), *Xalatan* (November 2006), *Thyrolar* (April 2005), *Kaletra* (November 2007), Lorazepam *Intensol* oral concentrate (June 2007), *Alkeran* (November 2004), *Sandostatin* (September 2005), *Sandostatin LAR* (March 2008), *Ethezyme* (March 2006), *Pegasys* (January 2008), *PegIntron* (April 2008), *Symlyn* (July 2008), *Promethegan* suppositories (June 2000), *Proparacaine* ophthalmic solution (July 2005), *Norvir* capsules (August 2008), *Veregen* (December 2007), *Rapamune*

(January 2008), *Forteo* (February 2008), *Tobi* (December 2007), *Viroptic* (October 2001), Diphtheria and Tetanus Toxoids, adsorbed (December 2005), *Decavac* (March 2004), *Tripedia* (March 2001), *Daptacel* (March 2008), *Pentacel* (June 2008), *Infanrix* (July 2003), *Pediarix* (March 2007), *Boostrix* (December 2005), *PedvaxHIB* (January 2001), *ActHib* (December 2005), *Comvax* (September 1996), *Havrix* (March 2008), *Vaqta* (May 2007), *Twinrix* (March 2007), *Engerix-B* (January 2007), *Recombivax* (May 2007), *Gardasil* (September 2008), *IPOL* (December 2005), *Afluria* (July 2008), *FluLaval* (July 2008), *Fluarix* (August 2008), *Fluvirin* (August 2008), *Fluzone* (August 2008), *FluMist* (July 2008), *M-M-R II* (May 2008), *ProQuad* (February 2008), *Menactra* (October 2007), *Menomune* (December 2005), *Prevnar* (December 2007), *Pneumovax 23* (August 2008), *RotaTeq* (April 2008), *Rotarix* (April 2008), *Varivax* (November 2006), *Vivotif* (August 2006), *Zostavax* (July 2007).

***** Room temperature stability determined on a case-by-case basis. Contact manufacturer and provide lot number and expiration date, length of time exposed to room temperature, maximum exposure temperature.**

§ Off-label information is not reviewed or approved by the FDA.

Users of this document are cautioned to use their own professional judgment and consult any other necessary or appropriate sources prior to making clinical judgments based on the content of this document. Our editors have researched the information with input from experts, government agencies, and national organizations. Information and Internet links in this article were current as of the date of publication.

Safeguarding Pharmaceutical Products During Power Outages or Flooding

In the event of a **power outage**, there are a number of steps to take to minimize damage to pharmaceutical products. If the power goes out, the time of the power outage should be noted. During the power outage, refrigerator and freezer doors should not be opened. For products which are stored in the freezer, if the power outage continues for more than 24 hours, the products should be removed from the freezer and packed in dry ice.

When the power is restored, the temperature in the refrigerator and freezer should be recorded, and monitored until the temperature has returned to the desired temperature. This allows the calculation of the duration of time the product was stored at a temperature outside of the product labeling and may enable calculations to be made by the product manufacturer, in consultation with FDA as necessary, about the continued potency of the involved products.³

When a **flood** is anticipated, if possible, pharmaceutical products should be moved out of areas which could accumulate flood water. If possible, elevate products stored on warehouse floors off the ground (e.g., on pallets). For those items on shelves, securely anchor the shelves to keep products dry. For refrigerated products, raise refrigerators onto platforms, if possible.³

If flooding occurs, any product which was not sealed in airtight, waterproof bags should be considered contaminated and should be discarded.³

Users of this document are cautioned to use their own professional judgment and consult any other necessary or appropriate sources prior to making clinical judgments based on the content of this document. Our editors have researched the information with input from experts, government agencies, and national organizations. Information and Internet links in this article were current as of the date of publication.

Project Leader in preparation of this Detail-Document: Neeta Bahal O'Mara, Pharm.D., BCPS

References

1. Cohen V, Jellinek SP, Teperikidis L, et al. Room-temperature stability of medications labeled for refrigeration. *Am J Health Syst Pharm* 2007;64:1711-5.
2. Welte MS. Vaccines ruined by poor refrigeration. *Washington Post*, December 4, 2007. <http://www.washingtonpost.com/wp-dyn/content/article/2007/12/04/AR2007120400983.html>. (Accessed December 5, 2007).
3. Food and Drug Administration. Impact of severe weather conditions on biological products. <http://www.fda.gov/cber/weatherimpact.htm>. (Accessed September 2, 2008).
4. Personal communication. Bethany Z. Dimaculangan, Pharm.D. Medical Information. Upsher-Smith Laboratories. Maple Grove, MN 55369. September 12, 2008.
5. Personal communication. Jacquelyn Collins. Medical Information. GlaxoSmithKline. Philadelphia, PA 19101. September 12, 2008.
6. Personal communication. Prachi Parmar, R.Ph. Medical Information. Pfizer. New York, NY 100017. September 12, 2008.
7. Personal communication. Deborah Bialy. Corporate Quality Product Support. Genentech. South San Francisco, CA 94080. September 12, 2008.
8. Personal communication. Leah Bernstein, Ph.D. Medical Information Department. Solvay Pharmaceuticals. Marletta, GA 30062. September 15, 2008.
9. Personal communication. Maleeha Ansari, Pharm.D. Global Medical Information. Schering Corporation. Kenilworth, NJ 07033. September 15, 2008.
10. Personal communication. Technical Product Information Department. BI Roxane. Columbus, OH 43216. September 15, 2008.
11. Personal communication. Michelle Kraus, Pharm.D., Medical Communications. KV Pharmaceutical. St. Louis, MO 63043. September 15, 2008.
12. Personal communication. Jessie Li, Pharm.D. Vaccine and Anti-infectives Team. GlaxoSmithKline. Philadelphia, PA 19101. September 12, 2008.
13. Personal communication. Gregory M. Susla, Pharm.D., FCCM. MedImmune. Gaithersburg, MD 20878. September 11, 2008.
14. Personal communication. Gloria Rios-Kretvix. Customer Communications Center. Centocor Ortho Biotech. Raritan, NJ 08869. September 23, 2008.
15. Personal communication. Diane Butner RN Scientific Affairs. Ortho-McNeil Janssen. Raritan, NJ 08869. September 23, 2008.
16. Personal communication. Amgen Canada. Mississauga, Ontario L5N 0A4. 866-502-6436. September 19, 2008.
17. Personal communication. Donna Clark. Information Specialist II. Professional Information Services. King Pharmaceuticals. Bristol, TN 37620. September 23, 2008.

Cite this Detail-Document as follows: *Stability of refrigerated and frozen drugs. Pharmacist's Letter/Prescriber's Letter 2008;24(10):241001.*

More . . .



Evidence and Advice You Can Trust...



3120 West March Lane, P.O. Box 8190, Stockton, CA 95208 ~ TEL (209) 472-2240 ~ FAX (209) 472-2249
Copyright © 2008 by Therapeutic Research Center – Posted with permission CR09312

Subscribers to *Pharmacist's Letter* and *Prescriber's Letter* can get *Detail-Documents*, like this one, on any topic covered in any issue by going to www.pharmacistsletter.com or www.prescribersletter.com